

梶川脳神経内科英文業績

(2011年まで)

- 1) **Nomura E, Kohriyama T, Yamaguchi S, Kajikawa H, Nakamura S**. Association between carotid atherosclerosis and hemostatic markers in patients with cerebral small artery disease. *Blood Coagul Fibrinolysis* 9: 55-62, 1998. (広島大学大学院脳神経内科との共同研究)
- 2) **Murata Y, Yamaguchi S, Kajikawa H, Yamamura K, Sumioka S, Nakamura S**. Relationship between the clinical manifestations, computed tomographic findings and the outcome in 80 patients with primary pontine hemorrhage. *J Neurol Sci* 167:107-111, 1999.
- 3) **Nomura E, Kohriyama T, Kozuka K, Kajikawa H, Nakamura S**. Sequential changes in von Willebrand factor and soluble thrombomodulin in acute ischemic stroke. *J Stroke Cerebrovasc Dis* 10: 257-264, 2001. (広島大学大学院脳神経内科との共同研究)
- 4) **Kozuka K, Kohriyama T, Nomura E, Ikeda J, Kajikawa H, Nakamura S**. Endothelial markers and adhesion molecules in acute ischemic stroke—sequential change and differences in stroke subtype. *Atherosclerosis* 161: 161-168, 2002. (広島大学大学院脳神経内科との共同研究)
- 5) **Nomura E, Kohriyama T, Kozuka K, Kajikawa H, Nakamura S, Matsumoto M**. Significance of serum soluble thrombomodulin level in acute cerebral infarction. *Eur J Neurol* 11: 329-334, 2004. (広島大学大学院脳神経内科との共同研究)
- 6) **Naka H, Nomura E, Wakabayashi S, Kajikawa H, Kohriyama T, Mimori Y, Nakamura S, Matsumoto M**. Frequency of Asymptomatic Microbleeds on T2*-Weighted MR Images of Patients with Recurrent Stroke: Association with Combination of Stroke Subtypes and Leukoaraiosis. *Am J Neuroradiol* 25:714-719, 2004.
- 7) **Naka H, Nomura E, Takahashi T, Wakabayashi S, Kajikawa H, Kohriyama T, Mimori Y, Matsumoto M**. Plasma total homocysteine levels are associated with advanced leukoaraiosis but not with asymptomatic microbleeds on T2*-weighted MRI in patients with stroke. *Eur J Neurol* 13: 261-265, 2006.

- 8) **Naka H, Nomura E, Takahashi T, Wakabayashi S, Mimori Y, Kajikawa H, Kohriyama T, Matsumoto M**. Combinations of the presence or absence of cerebral microbleeds and advanced white matter hyperintensity as predictors of subsequent stroke types. *Am J Neuroradiol* 27:830-835, 2006.
- 9) **Ueno H, Naka H, Ohshita T, Kondo K, Nomura E, Ohtsuki T, Kohriyama T, Wakabayashi S, Matsumoto M**. Association between cerebral microbleeds on T2*-Weighted MR images and recurrent hemorrhagic stroke in patients treated with warfarin following ischemic stroke. *Am J Neuroradiol* 29: 1483-1486, 2008. (広島大学大学院脳神経内科との共同研究)
- 10) **Lovelock CE, Cordonnier C, Naka H, Salmon R Al-S, Sudlow CLM, Edinburgh Stroke Study Group, Sorimachi T, Werring DJ, Gregoire SM, Imaizumi T., Lee S-H, Briley D., Rothwell PM**. Antithrombotic drug use, cerebral microbleeds, and intracerebral hemorrhage: A systemic review of published and unpublished studies. *Stroke* 41:1222-1228, 2010. (国際共同研究)
- 11) **Sueda Y, Naka H, Ohtsuki T, Kono T, Aoki S, Ohshita T, Nomura E, Wakabayashi S, Kohriyama T, Matsumoto M**. Positional Relationship between Recurrent Intracerebral Hemorrhage/Lacunar Infarction and Previously Detected Microbleeds. *Am J Neuroradiol* 31:1498-1503, 2010. (広島大学大学院脳神経内科との共同研究)